

Functional Reach and Dynamic Balance of Healthy 11-Year-Old Children

Katerina Kirilova, Katerina Stambolieva, Plamen Gatev

Institute of Neurobiology, Bulgarian Academy of Sciences



INTRODUCTION

Standing balance depends on the adequate integration of sensory information and the support area [1]. Functional reach (FR) test [2] is applied for dynamic balance assessment of different age groups. Three strategies of FR have been reported: ankle, hip and mixed [3], defined by the changes in the ankle and hip joints during reach.

AIM


To compare the reaching strategies and dynamic balance of healthy 11-year-old children and adult controls and to assess children's ability to adapt to different task conditions.

SUBJECTS

Eighteen children – boys and girls (mean age 11 ± 0.2 years) and 19 adults (mean age 32.4 ± 4.9 years)

EXPERIMENTAL SETUP & PROCEDURE

- Two trials of FR per task condition
- Four sensory conditions:
 - eyes open (EO) – all sensory information available
 - eyes closed (EC) – no visual information
 - eyes open, head extended (EOHE) – altered vestibular information
 - eyes closed, head extended (ECHE) – no visual and altered vestibular information
- Two types of support area (SA):



- Pedobarographic platform Tekscan: recording center of pressure (COP) sway, 30 fps, 30s.
- GoPro Hero 5 Black camera: video recording of kinematic data, 1280x720 px, 60 fps.
- Specialized software:
 - Kinovea v0.8.26 for automatic object tracking
 - custom-made programs in MatLab for 2D kinetic and kinematic analysis [4].

EVALUATED MEASURES

- Medio-lateral sway of COP
- FR length
- FR normalized by height
- Changes in the hip and ankle joint angles before and during FR

RESULTS AND DISCUSSION

We defined two strategies – hip and ankle (see photos below) by the differences in the ankle and hip joints during the maximal forward reach and the initial body position. Almost all subjects adopted the hip strategy (Table 1). Subjects were rather consistent in their choice of reaching strategy in all task conditions (Table 2). As few people adopted the ankle strategy, FR and medio-lateral COP sway were further evaluated only for the subjects with hip strategy.

HIP STRATEGY

Hip angle considerably decreases. Ankle angle increases.



Hip angle changes during FR


		Normal support				Narrow support			
		EO	EC	EOHE	ECHE	EO	EC	EOHE	ECHE
Adults n = 16	mean	-22.9	-22.8	-22.4	-22.8	-22.0	-21.5	-21.3	-21.6
	min	-43.2	-40.1	-42.9	-42.4	-37.5	-39.6	-36.9	-42.8
	max	-8.0	-7.2	-5.2	-4.3	-6.8	-7.2	-4.1	-5.6
	SEM	2.135	2.239	2.317	2.119	1.850	1.989	1.904	1.976
Children n = 13	mean	-26.0	-24.3	-26.4	-25.0	-26.8	-24.2	-24.5	-23.5
	min	-49.5	-42.5	-53.2	-42.3	-50.0	-54.3	-54.1	-44.2
	max	-10.8	-8.8	-10.2	-12.0	-16.0	-11.0	-12.0	-11.9
	SEM	3.120	2.572	3.629	2.807	2.699	2.779	2.966	2.307

Ankle angle changes during FR

		Normal support				Narrow support			
		EO	EC	EOHE	ECHE	EO	EC	EOHE	ECHE
Adults n = 16	mean	8.4	8.6	8.6	8.6	9.6	10.0	9.9	9.7
	min	4.4	5.4	5.0	4.1	5.7	5.5	5.8	5.8
	max	13.7	12.4	11.5	11.4	14.3	16.6	13.0	12.6
	SEM	0.649	0.588	0.570	0.541	0.641	0.743	0.533	0.569
Children n = 13	mean	10.0	10.4	8.9	9.3	12.1	11.5	11.6	10.9
	min	6.2	4.3	5.0	4.5	6.1	6.6	5.4	5.8
	max	17.1	19.5	15.0	14.1	20.6	19.6	19.5	16.0
	SEM	0.842	0.940	0.715	0.780	0.971	1.082	1.127	0.837

ANKLE STRATEGY

Smaller changes in hip angle. Ankle angle decreases.



Hip angle changes during FR


		Normal support				Narrow support			
		EO	EC	EOHE	ECHE	EO	EC	EOHE	ECHE
Adults n = 3	mean	-3.0	-3.4	-2.0	-3.4	-2.6	-2.6	-1.4	-3.5
	min	-6.5	-6.9	-5.5	-8.2	-5.3	-5.0	-3.9	-6.5
	max	1.5	2.3	2.3	2.4	1.9	1.6	1.8	0.8
	SEM	0.937	1.135	0.908	1.225	0.891	0.841	0.673	0.871
Children n = 5	mean	1.1	0.3	0.7	-1.3	0.3	0.7	1.2	-0.8
	min	-2.7	-1.4	-1.8	-7.3	-2.4	-0.9	-0.4	-2.6
	max	6.5	3.1	2.3	2.0	2.2	2.3	3.5	0.7
	SEM	1.499	0.811	0.694	1.583	0.797	0.579	0.749	0.598

Ankle angle changes during FR


		Normal support				Narrow support			
		EO	EC	EOHE	ECHE	EO	EC	EOHE	ECHE
Adults n = 3	mean	-0.6	-0.6	-1.4	0.2	-0.3	-0.4	-1.4	0.5
	min	-1.6	-2.0	-3.6	-1.0	-2.4	-2.1	-2.9	-1.1
	max	0.8	1.4	1.5	0.9	1.8	1.8	1.1	2.4
	SEM	0.294	0.399	0.599	0.252	0.481	0.460	0.502	0.411
Children n = 5	mean	-2.3	-2.6	-2.6	-3.2	-2.1	-3.0	-2.9	-1.8
	min	-5.5	-3.9	-3.9	-4.2	-4.1	-4.3	-4.8	-3.4
	max	1.8	-1.2	-0.7	-2.2	0.0	-2.3	0.2	0.0
	SEM	1.182	0.612	0.591	0.377	0.655	0.339	0.864	0.713

Table 2. Descriptive statistics of the changes in the hip (up) and ankle (down) joint angles before and during FR in all task conditions for hip (left) and ankle (right) strategies.

Functional Reach Normalized by Body Height



Medio-Lateral Sway of COP



Mean values of FR normalized by height (left) and M-L COP sway of the groups with hip strategy on normal and narrow support area (SA) in the four sensory conditions (see text). * - differences between sensory conditions; ^ - differences between series on normal and narrow support, $p < 0.05$ (paired t-test).


Sensory conflict diminished FR length similarly in both age groups. Task performance was not affected by the narrow support. Children had shorter FR length in general, but when normalized by body height, their performance didn't differ significantly from the adult controls. Children's lateral sway was slightly greater in general. Sensory conflict and/or narrowing the support increased the lateral COP sway in both age groups. The results are in line with previous studies [5, 6] that also found adult-like postural behavior in children. However, children's dynamic balance seems to be more visually dependent.

CONCLUSIONS

The sensory conflict and narrow support deteriorate the dynamic standing balance of both children and adults. The 11-year-old children showed adult-like adaptation to the different task constraints.

Table 1: Number of people of each group and adopted FR strategies.

ACKNOWLEDGEMENTS



The study was funded by Project Grants DTK 02/60 with the National Science Fund and DFNP 17-55 with the Ministry of Education and Science, Republic of Bulgaria. www.cleverstance.com

REFERENCES

- Day B, Steiger M, Thomson P, Marsden C. Effect of vision and stance width on human body motion when standing: implications for afferent control of lateral sway. *J. Physiology*. 1993, vol. 469, p. 479-499.
- Duncan P, Weiner D, Chandler J, Studenski S. Functional reach: A new clinical measure of balance. *J Gerontol*. 1990, vol. 45(6), p. M192-197.
- Liao C, Lin S. Effects of different movement strategies on forward reach distance. *Gait Posture*. 2008 Jul; vol. 28(1), p. 16-23.
- Kirilova K, Gatev P. 2018. A complex semi-automatic method for kinetic and two-dimensional kinematic motion analysis for posture and movement investigation. *Int. J. Bioautomation*. 2018, vol. 22(1), p. 57-64.
- Mallau S, Vaugoyeau M, Assaiante, C. Postural strategies and sensory integration: no turning point between childhood and adolescence. *PLoS One*. 2010, vol. 5(9), e13078.
- Hatzitaki V, Zisi V, Kollias I, Kioumourtzoglou E. Perceptual-motor contributions to static and dynamic balance control in children. *J. Mot. Behav*. 2002, vol. 34(2), p. 161-170.